

Short Commentary

Effect of Chronic Circadian Disorder during Pregnancy on Progeny Mice**Zhang P^{*1}**

¹Key Laboratory of Medical Electrophysiology of Ministry of Education and Medical Electrophysiological Key Laboratory of Sichuan Province, Institute of Cardiovascular Research at Southwest Medical University, Luzhou 646000, China.

**Corresponding author:* Zhang P, Key Laboratory of Medical Electrophysiology of Ministry of Education and Medical Electrophysiological Key Laboratory of Sichuan Province, Institute of Cardiovascular Research at Southwest Medical University, Luzhou 646000, China; Email: Peng_Zhang@brown.edu

Received Date: 09-11-2018

Accepted Date: 09-18-2018

Published Date: 12-10-2018

Copyright: © 2018 Zhang P

Circadian rhythms are very important to the reproduction [1, 2], development [3, 4] and physiological activities of autonomic nerve innervation of mammals and can even be said to be: Time is everything. Here is a short communication about 'Environmental perturbation of the circadian clock during pregnancy leads to transgenerational mood disorder-like behaviors in mice' which we published in 2017. At the same time, some interesting scientific questions will be raised up, which are also the substance of our further research.

In the study of the relationship between circadian rhythm and various diseases, mental disease is not a popular direction. Some studies have found a strong link between the rhythm of master clock, SCN, and a variety of mental disorders, like bipolar and depression disorder [5-8]. A large proportion of patients with major depression disorder, anxiety and bipolar disorder are accompanied by varying degrees of sleep-arousal disorder [9,10]. Even the first symptom of some patients with mental illness is sleep-arousal disorder [11]. But it is difficult to identify the causal relationship between mental disorder and circadian disorder. Although our research is not sufficient to clarify the relationship between the two. Our findings may have

some implications for this: they may be causally related to each other. Many articles reported that chronic restraint stress or chronic unpredictable stimulation in pregnancy can cause abnormal behavior of offspring mice or rats similar to mental disorders [12-14]. On the other side, the abnormal circadian rhythms were found in the detection of SCN circadian rhythm in MDD model rats and mice [15,16]. Based on these backgrounds, our initial assumption was: would it be possible to induce depression or anxiety in mice if they were stimulated by a single rhythmic disorder? However, it was found that frequent switching circadian rhythms did not induce significant behavioral abnormalities in adult mice (forced swimming and sugar water preference experiments). So, what about the effect on the offspring? Then we designed frequent circadian rhythms of maternal mice during pregnancy. The results showed that offspring mice had subculture abnormal behavior (at least 1 generation). At this point, we found that rhythm disorder can be the cause of mental disorders.

Embryonic development is a rigorous and orderly stage of space-time development the timeline is a crucial rule in this development, and once the schedule is broken, it can have a fatal effect on the embryo in which organisms

undergo programmed changes in fixed time and space. We can imagine that mammalian embryos cannot directly perceive light signals in the environment of the mother undefined womb, but only through the mother undefined neural fluid to regulate the passive acceptance of the timing signal. Although we know that the programmed spatiotemporal expression patterns of embryonic development are determined by genes, However, abnormal maternal changes, such as changes in the levels of nerve, body fluids and hormones caused by abnormal rest, feeding time, and the state of psych sperm, are both stress states for both embryos and embryos. This stress state is likely to lead to an increase in the incidence of gene mutations and the effect of epigenetic levels in embryos. Once this effect occurs, it may involve not only a single organ, but also a wide range of multiple systems. Therefore, we confirmed that the difference protein did involve multiple systems and biological functions after the protein differential spectrum was detected in the hypothalamus of children with chronic rhythm disorder during pregnancy. We tested the expression of MC4R, one of the differential proteins. And it was found that there was not only significant difference between NS and CCD mice, but also has gender difference in CCD-F1 mice. This has aroused our concern about the cardiovascular system and sexual development of the offspring after CCD.

MC4R is, which is G protein-coupled receptors that stimulate adenylate cyclase, one of five melanocortin receptors (MCRs) (MC1R-MC5R) [17-20]. MC4Rs are mainly present in the central nervous system (CNS). Melanocortins responds to stress [21], and melanocortins regulate several physiological functions, including body weight, sexual behavior, and cardiovascular function [22-29]. The research indicated that α -MSH microinjections into the nucleus ambiguous exert excitatory effects on parasympathetic preganglionic nucleus ambiguous neurons via MC4Rs, leading to bradycardic responses. [30]. We spontaneously thought that do CCD-F1 mice has abnormal phenotypes in cardiovascular activity? It there a sexual dimorphism?

Therefore, we observed the CCD-F1 generation mice of different genders. According to the experimental results obtained at present, CCD-F1-female has obvious abnormal puberty development, which is characterized

by delayed vaginal opening time and disorder of estrous cycle. The phenotypes of CCD-F1-male were mainly found in the increased susceptibility to arrhythmias. Under the same dose of drug induction, fatal arrhythmias were more likely to occur (ventricular fibrillation, mortality rate was much higher than that of non-stressed F1 generation). We will continue to study these two interesting phenomena. We hope that our results can further understand the regulation of daily rhythm on the development and function of different tissues and organs.

References

1. Boden MJ, Varcoe TJ, Kennaway DJ. Circadian regulation of reproduction: from gamete to offspring. *Prog Biophys Mol Biol.* 2013; 113(3): 387-97.
2. Beymer M, Henningsen J, Bahougne T, Simonneaux V. The role of kisspeptin and RFRP in the circadian control of female reproduction. *Mol Cell Endocrinol.* 2016; 438: 89-99.
3. Landgraf D1, Achten C, Dallmann F, Oster H. Embryonic development and maternal regulation of murine circadian clock function. *Chronobiol Int.* 2015; 32(3): 416-27.
4. Vallone D1, Lahiri K, Dickmeis T, Foulkes NS. Start the clock! Circadian rhythms and development. *Dev Dyn.* 2007; 236(1): 142-55.
5. Wu X, Balesar R, Lu J, Farajnia S, Zhu Q, et al. Increased glutamic acid decarboxylase expression in the hypothalamic suprachiasmatic nucleus in depression. *Brain Struct Funct.* 2017; 222(9): 4079-4088.
6. Landgraf D, Joiner WJ, McCarthy MJ, Kiessling S, Barandas R, et al. The mood stabilizer valproic acid opposes the effects of dopamine on circadian rhythms. *Neuropharmacology.* 2016; 107: 262-270.
7. Logan RW, Edgar N, Gillman AG, Hoffman D, Zhu X, et al. Chronic Stress Induces Brain Region-Specific

- Alterations of Molecular Rhythms that Correlate with Depression-like Behavior in Mice. *Biol Psychiatry*. 2015; 78(4): 249-58.
8. Nagy AD, Iwamoto A, Kawai M, Goda R, Matsuo H, et al. Melatonin adjusts the expression pattern of clock genes in the suprachiasmatic nucleus and induces antidepressant-like effect in a mouse model of seasonal affective disorder. *Chronobiol Int*. 2015; 32(4): 447-57.
 9. Tsujimoto T, Yamada N, Shimoda K, Hanada K, Takahashi S. Circadian rhythms in depression. Part II: Circadian rhythms in inpatients with various mental disorders. *J Affect Disord* 1990; 18(3): 199-210.
 10. Monteleone P, Martiadis V, Maj M. Circadian rhythms and treatment implications in depression. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011; 35(7): 1569-74.
 11. Cameron OG, Lee MA, Kotun J, McPhee KM. Circadian symptom fluctuations in people with anxiety disorders. *J Affect Disord*. 1986; 11(3): 213-8.
 12. Kiryanova V, Smith VM, Antle MC, Dyck RH. Behavior of Adult 5-HT1A Receptor Knockout Mice Exposed to Stress During Prenatal Development. *Neuroscience*. 2018; 371: 16-28.
 13. Kiryanova V, Smith VM, Dyck RH, Antle MC. Circadian behavior of adult mice exposed to stress and fluoxetine during development. *Psychopharmacology (Berl)*. 2017; 234(5): 793-804.
 14. Chung S, Son GH, Park SH, Park E, Lee KH, et al. Differential adaptive responses to chronic stress of maternally stressed male mice offspring. *Endocrinology*. 2005; 146(7): 3202-10.
 15. Griesauer I, Diao W, Ronovsky M, Elbau I, Sartori S, et al. Circadian abnormalities in a mouse model of high trait anxiety and depression. *Ann Med*. 2014; 46(3): 148-54.
 16. Tapia-Osorio A, Salgado-Delgado R, Angeles-Castellanos M, Escobar C. Disruption of circadian rhythms due to chronic constant light leads to depressive and anxiety-like behaviors in the rat. *Behav Brain Res*. 2013; 252: 1-9.
 17. Hadley ME, Hruby VJ, Jiang J, Sharma SD, Fink JL, et al. Melanocortin receptors: identification and characterization by melanotropic peptide agonists and antagonists. *Pigment Cell Res*. 1996; 9(5): 213-234.
 18. Adan RA, Gispen WH. Melanocortins and the brain: from effects via receptors to targets. *Eur J Pharmacol*. 2000; 405: 13-24.
 19. Kishi T, Aschkenasi CJ, Lee CE, Mountjoy KG, Saper CB, Elmquist JK. Expression of melanocortin 4 receptor mRNA in the central nervous system of the rat. *J Comp Neurol*. 457: 2003; 457(3): 213-235.
 20. Wikberg JES, Mutulis F. Targeting melanocortin receptors: an approach to treat weight disorders and sexual dysfunction. *Nat Rev Drug Discov*. 2008; 7(4): 307-323.
 21. Liu J, Garza JC, Truong HV, Henschel J, Zhang W, Lu XY. The melanocortinergic pathway is rapidly recruited by emotional stress and contributes to stress-induced anorexia and anxiety-like behavior. *Endocrinology*. 2007; 148(11): 5531-5540.
 22. Brown S, Chitravanshi VC, Sapru HN. Cardiovascular actions of adrenocorticotropin into the nucleus tractus solitarius of the rat. *Neuroscience*. 2006; 143(3): 863-874.
 23. Hill C, Dunbar JC. The effects of acute and chronic alpha melanocyte stimulating hormone (alpha-MSH) on cardiovascular dynamics in conscious rats. *Peptides*. 2002; 23(9):1625-30: 1625-1630.

-
24. Humphreys MH. Cardiovascular and renal actions of melanocyte-stimulating hormone peptides. *Curr Opin Nephrol Hypertens.* 2007; 16 (1): 32 –38.
25. Kawabe T , Chitravanshi VC, Kawabe K, Sapru HN. Cardiovascular effects of adrenocorticotropin microinjections into the rostral ventrolateral medullary pressor area of the rat. *Brain Res.* 2006; 1102(1): 117 –126.
26. Ni XP, Butler AA, Cone RD, Humphreys MH. Central receptors mediating the cardiovascular actions of melanocyte stimulating hormones. *J Hypertens.* 2006; 24(11): 2239 –2246.
27. Ramaekers D, Beckers F, Demeulemeester H, Bert C, Denef C, et al. Effects of melanocortins on cardiovascular regulation in rats. *Clin Exp Pharmacol Physiol.* 2002; 29(7): 549 –558.
28. Rinne P , Harjunpaa J, Scheinin M, Savontaus E. Blood pressure regulation and cardiac autonomic control in mice overexpressing alpha- and gamma-melanocyte stimulating hormone. *Peptides.* 2008; 29(11): 1943 –1952.
29. Versteeg DH, Bergen PV, Adan RA, De Wildt DJ. Melanocortins and cardiovascular regulation. *Eur J Pharmacol.* 1998; 360(1):1-14.
30. Chitravanshi VC1, Bhatt S, Sapru HN. Microinjections of alpha-melanocyte stimulating hormone into the nucleus ambiguus of the rat elicit vagally mediated bradycardia. *Am J Physiol Regul Integr Comp Physiol.* 2009; 296(5): R1402-11.